



REST is a hypoxia-responsive transcriptional repressor

Cavadas, M. A. S., Mesnieres, M., Crifo, B., Manresa, M. C., Selfridge, A. C., Keogh, C. E., Fabian, Z., Scholz, C. C., Nolan, K. A., Rocha, L. M. A., Tambuwala, M., Brown, S., Wdowicz, A., Corbett, D., Murphy, K. J., Godson, C., Cummins, E. P., Taylor, C. T., & Cheong, A. (2016). REST is a hypoxia-responsive transcriptional repressor. *Scientific Reports*, 6, [31355 (2016)]. <https://doi.org/10.1038/srep31355>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Scientific Reports

Publication Status:
Published (in print/issue): 17/08/2016

DOI:
[10.1038/srep31355](https://doi.org/10.1038/srep31355)

Document Version
Author Accepted version

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

REST is a hypoxia-responsive transcriptional repressor.

Miguel A. S. Cavadas^{1,2,3}, Marion Mesnieres², Bianca Crifo², Mario C. Manresa², Andrew C. Selfridge², Ciara E Keogh², Zsolt Fabian², Carsten C. Scholz^{1,2,10}, Karen A. Nolan^{4,10}, Liliane M. A. Rocha⁵, Murtaza M. Tambuwala⁶, Stuart Brown⁷, Anita Wdowicz⁸, Danielle Corbett⁸, Keith J. Murphy⁸, Catherine Godson⁴, Eoin P. Cummins², Cormac T. Taylor^{1,2,*} and Alex Cheong^{1,2,9,*}

¹ Systems Biology Ireland, University College Dublin, Dublin 4, Ireland.

² Conway Institute of Biomolecular and Biomedical Research, School of Medicine and Medical Sciences, University College Dublin, Dublin 4, Ireland.

³ Instituto Gulbenkian de Ciência, Rua da Quinta Grande, 2780-156 Oeiras, Portugal.

⁴ Diabetes Complications Research Centre, School of Medicine and Medical Sciences, University College Dublin, Dublin 4, Ireland.

⁵ Faculdade de Medicina, Universidade de Lisboa, 1649-028 Lisbon, Portugal.

⁶ School of Pharmacy and Pharmaceutical Sciences, University of Ulster, Coleraine, Co. Londonderry, BT52 1SA, Northern Ireland, UK.

⁷ Center for Health Informatics and Bioinformatics, New York University School of Medicine, New York, NY 10016, USA.

⁸ Neurotherapeutics Research Group, UCD School of Biomolecular and Biomedical Science, Conway Institute, University College Dublin, Belfield, Dublin 4, Ireland.

⁹ Life and Health Sciences, Aston University, Birmingham , B4 7ET, UK.

¹⁰ Institute of Physiology and Zurich Centre for Integrative Human Physiology, University of Zurich, Zurich, Switzerland.

*These authors contributed equally to this work

Correspondence: Dr Alex Cheong, Life and Health Science, Aston University, Birmingham B4 7ET, +44-1212043420, email: a.cheong@aston.ac.uk

Abstract

Cellular exposure to hypoxia results in altered gene expression in a range of physiologic and pathophysiologic states. Discrete cohorts of genes can be either up- or down-regulated in response to hypoxia. While the Hypoxia-Inducible Factor (HIF) is the primary driver of hypoxia-induced adaptive gene expression, less is known about the signalling mechanisms regulating hypoxia-dependent gene repression. Using RNA-seq, we demonstrate that equivalent numbers of genes are induced and repressed in human embryonic kidney (HEK293) cells. We demonstrate that nuclear localization of the Repressor Element 1-Silencing Transcription factor (REST) is induced in hypoxia and that REST is responsible for regulating approximately 20% of the hypoxia-repressed genes. Using chromatin immunoprecipitation assays we demonstrate that REST-dependent gene repression is at least in part mediated by direct binding to the promoters of target genes. Based on these data, we propose that REST is a key mediator of gene repression in hypoxia .

Keywords: Hypoxia, Repressor, REST, transcription.

Introduction

Hypoxia is a feature of a range of physiological and pathophysiological conditions including embryonic development, exercise, cancer, ischemia and inflammation ¹. Throughout evolution, adaptive pathways have developed to help an organism cope with hypoxia. The best-described transcriptional adaptive response in cells is mediated by the hypoxia inducible factor (HIF) signalling pathway, which up-regulates genes which restore oxygen and energy homeostasis ²⁻⁴. In normoxia, HIF α is hydroxylated by the prolyl-hydroxylase domain (PHD) family of dioxygenases targeting it for ubiquitylation by the von Hippel Lindau E3 ligase complex and subsequent proteosomal degradation ⁵. This process is reversed in hypoxia and HIF α is stabilized, dimerises with HIF β and binds to hypoxia response elements (HRE) in the regulatory regions of target genes ⁶. HIF drives an adaptive response to hypoxia by promoting the expression of genes that regulate erythropoiesis, angiogenesis and glycolysis ⁶. However in cancer, HIF signalling can be maladaptive and contribute to tumour survival ⁷.

Microarray studies of mammalian cells exposed to hypoxia have shown that approximately the same numbers of genes are repressed and induced ^{2,3,8}. While HIF has been described as a master regulator of gene expression in hypoxia, a significant number of other transcription factors are also hypoxia-sensitive and control HIF-independent gene expression ⁹. Microarray data combined with siRNA against HIF-1/-2 indicate that there are HIF-dependent and HIF-independent genes which are differentially expressed in hypoxia ². ChIP-seq and microarray data indicate that while HIF is enriched in the promoters of genes induced in a HIF-dependent way, it is not enriched in the promoters of genes that are repressed in hypoxia, thus indicating the presence of HIF-independent and/or indirectly HIF-dependent mechanisms governing gene repression in hypoxia ¹⁰. While HIF is well known to regulate gene induction ²⁻⁴, the mechanisms underpinning transcriptional repression of genes in hypoxia remain poorly understood and is the topic of the current study ^{8,11}. Transcriptional repressors are a large and diverse group of proteins ¹². Mechanistically repressors can act by inhibiting the basal transcription machinery, ablation of activator function and remodelling of chromatin. They can be further classified into DNA-binding proteins (Class I) like Krüppel zinc fingers, proteins that bind DNA-binding proteins (Class II) such as the DNA-methyltransferase Dnmt3, or proteins that target activators, reducing their activity, such as I κ B that sequesters NF κ B in the cytosol ¹².

The Repressor Element 1-Silencing Transcription Factor (REST) is a C2H2- or Krüppel-type zinc finger, one of the largest classes of transcription factors in humans ¹³. It binds to the 21 base pair Repressor Element 1 (RE1) on the promoter of target genes and inhibits transcription by regulating

chromatin structure or by inhibiting the basal transcriptional machinery¹⁴. Proteosomal REST degradation is induced during neuronal differentiation, resulting in the promotion of the expression of the genes which confer the unique neuronal phenotype^{14,15}. Thus, REST was initially regarded as a master regulator of neurogenesis and as the first example of a vertebrate transcription silencer protein that regulates a large repertoire of cell type-specific genes¹⁶. However, REST has since being shown to be implicated in the regulation of non-neuronal biological processes in a variety of cell types, including cardiac myocytes, immune, vascular and tumour cells^{15,17-19}. In the aging human brain, REST is induced to protect against oxidative stress and brain neurodegeneration²⁰. Of particular interest for our study, ischemia and/or oxygen-glucose deprivation have previously been shown to induce REST nuclear protein and mRNA^{21,22} and to modulate the expression of target genes²¹. Furthermore, we have recently shown that REST represses the HIF-1 α mRNA expression and contributes to the resolution of the HIF response during prolonged hypoxia²³. This led us to explore the global response of REST to cellular hypoxia, with the aim of developing our understanding of the signalling mechanisms underpinning hypoxia-dependent transcriptional repression.

Using an unbiased, high-throughput approach combined with biochemical analysis, we demonstrate that REST accumulates in the nucleus of cells exposed to hypoxia and acts as a key repressor of the hypoxic transcriptome, regulating approximately 20% of the hypoxia-repressed genes. Furthermore, hypoxia leads to a change in the target gene repertoire of REST, from the repression of neuronal genes in normoxic/basal conditions, to the repression of metabolic, cell cycle and proliferation related genes in hypoxia. Together these findings indicate a previously unknown key role for the tumour suppressor REST in hypoxic signalling.

Results

High-throughput analysis of the transcriptional responses to hypoxia.

Previous microarray-based transcriptomic analysis of changes in global mRNA expression in response to hypoxia revealed that down-regulated genes reach their maximal repression following prolonged hypoxic exposure, while induced transcripts generally reach their maximum induction at earlier time-points⁸. Here we used human embryonic kidney (HEK293) cells exposed to normoxia or hypoxia for 24 hours as a cell model to investigate the processes associated with gene repression. High-throughput sequencing of poly(A)+ RNA (RNA-Seq) was performed on samples collected and transcript analysis revealed the presence of almost 2000 genes that were differentially expressed in hypoxia, with similar numbers of genes being induced (green; 851) and repressed (red; 1013) (Figure 1A). These data are quantitatively consistent with previously published transcriptomic studies of human cells exposed to hypoxia^{2,3,8}. Importantly, cohorts of well-characterized hypoxia-induced genes were up regulated including *LDHA*, *HK2*, *SLC2A1*, *EGLN1* and *BNIP3* (Figure 1B). Furthermore, a number of previously characterized hypoxia-repressed genes were present in the down-regulated cohort including *RRS1* (involved in ribosome biogenesis) and *MTHFD1* (involved in *de novo* purine synthesis) (Figure 1B)^{3,8}. Therefore, hypoxia had comparable effects on global gene induction and repression in HEK293 cells.

Using PANTHER ontological analysis, we classified hypoxia-induced and repressed genes according to gene ontology (<http://www.pantherdb.org/>). Our analysis revealed that hypoxia-induced and repressed genes could be associated with either similar or distinct processes (Figure 1C, Figures 1D, S1). For example, the glycolytic pathway, known previously to be induced in response to hypoxia¹, was identified only on the list of hypoxia-induced metabolic pathways (Figures 1C, S1G) while genes encoding proteins associated with intercellular junctions were identified only in the hypoxia-repressed gene cohort (Figure S1F). Although both induced and repressed gene cohorts included genes involved in the regulation of transcription, metabolism and development, only the repressed cohort contained an overrepresentation of genes involved in mRNA processing and splicing (Figures 1D). Taken together, these results indicate that hypoxia alters the expression of an equivalent number of increased and decreased genes which have both common and distinct roles in regulating cell responses.

REST is a hypoxia sensitive transcription factor

We next focused on possible molecular mechanisms underpinning the repression of gene transcription in hypoxia. REST was initially identified as a regulator of neuronal gene expression,

but was subsequently shown to have non-neuronal roles¹⁷⁻¹⁹. Using publically available microarray datasets, we confirm that REST is extensively expressed in multiple tissue types and cell lines, being highly expressed in HEK293 cells used in this study (Figure 2A and S2). HEK293 cells have been instrumental for the discovery of various aspects of REST biology: transcriptional networks²⁴, phosphorylation and the proteasome system²⁵ and regulation of HIF-1 α in hypoxia²³. Similar levels of REST could be detected in the nucleus and cytoplasm of normoxic HEK293 cells (Figure S3). In response to hypoxia, there was a more pronounced increase in REST levels in the nuclear (Figures 2B, C) than in either the cytoplasmic (Figures 2D, E) or whole cell extracts (Figure 2F,G), suggesting nuclear accumulation as the major response of REST to hypoxia. This nuclear accumulation was also observed in MCF10A (Figure S4). We investigated the effect of re-oxygenation on nuclear REST levels, and found that accumulation in hypoxia was reversible within one hour of re-oxygenation (Figure 2H), strongly supporting an oxygen-regulated post-translational control mechanism of REST cellular localization from the cytoplasm to the nucleus and back.

The increase in nuclear REST protein was associated with an increase in REST-dependent repressor activity, as measured using by a repressor element-1 (RE1)-luciferase assay developed to measure REST transcriptional activity (Figures 2I and S5). REST mRNA expression during hypoxic exposure was modestly and transiently increased, and was statistically significant only at the 2 hours time point (Figure 2J).

To assess the effect of hypoxia on REST protein stability, HEK293 cells were exposed to 24 hours of either normoxia (21 % oxygen) or hypoxia (1 % oxygen), followed by cycloheximide treatment in normoxia and protein stability was monitored by western blot. REST displayed a long half-life that was not increased by hypoxia (Figures S6A, B). In contrast, HIF-1 α in the same cells exhibited a short half-life, with hypoxia-stabilized HIF-1 α being completely degraded within 1 hour of culture in normoxia (Figure S6C). β -Actin exhibited a long half-life that was also not affected by hypoxia (Figure S6D).

Taken together, these results suggest that hypoxia leads to REST nuclear translocation, through a post-translational oxygen regulated mechanism, with a modest increase in transcription and, unlike HIF, independent of altered protein stability.

Role of REST in basal gene expression in hypoxia

The observation that nuclear REST expression and activity are increased led us to hypothesize that REST may play a broad role in regulating gene repression in hypoxia. To address this, we performed RNA-Seq analysis of hypoxic cells in the presence of scrambled or REST-specific siRNA (Figure 3A). An overview of the different steps performed to identify differentially expressed genes is shown in Figure S7A. All quality control checks are shown in Figures S7B-E and S8. As described earlier (Figure 1B), a cohort of known hypoxia-inducible genes served as positive experimental controls. REST-RNAi treatment led to a decrease in REST mRNA (Figures 3B and S9A) and protein expression (Figures S9B, S9C) and a resultant increase in a number of previously identified REST target genes^{24,26} in 21% oxygen (Figure 3C), thus validating the effectiveness of the REST-RNAi knockdown.

Among the genes whose basal expression was reduced in hypoxia (1013 genes represented by blue and red dots in Figure 3D), 201 (~20%) were found to be REST-Dependent (red dots in Figure 3D), i.e. they were not significantly repressed by hypoxia when REST was silenced (Figure 3E, Supplemental Table S1). On the other hand, 812 genes repressed in hypoxia were found to be REST-Independent, suggesting alternative gene regulation (Supplemental Table S1). An ontological and clustering analysis of the hypoxia-repressed genes showed that metabolic processes and other energy demanding processes including transcription, proliferation and cell cycle were heavily over-represented in the cohort repressed in a hypoxia and REST-Dependent manner (Figure 3F and Supplemental Table S2). Analysis of cell death in hypoxic cells treated with RNAi targeting REST did not reveal any effect of REST on cell death (Figure S10). Altogether, these data indicate that REST is an important transcriptional repressor in hypoxia, whose major function is beyond the control of cell survival in hypoxia.

REST regulates specific gene cohorts in normoxia and hypoxia by direct binding to gene promoters

Comparison of the REST-dependent genes repressed in hypoxia to the REST-Dependent genes repressed in normoxia revealed a limited overlap (5 genes, Figure 4A and Supplemental Table S3). Ontological analysis revealed well characterised neuronal genes regulated by REST in normoxia (Figures 4A black circle and 4B). A different subset of genes were found for REST-repressed genes in hypoxia (Figures 4A red circle and 4C), with ontological analysis showing an enrichment for metabolic and cell adhesion processes (Figure 4A). **Among the 10 genes most sensitive to REST regulation in hypoxia (Figure 4C), *RAB3C* (encoding for RAB3C GTPase), is predicted to be regulated by REST in normoxia and hypoxia, while *CYP1B1* (encoding for cytochrome P450 1B1) is predicted to be regulated only in hypoxia. Using qRT-PCR we tested and validated these**

findings (Figures 4D, E). Together, these results indicate that hypoxia leads to the recruitment of REST to a largely different set of genes from those regulated by REST in normoxia. We tested a further group of 6 genes for their regulation by REST in hypoxia. While we observed a similar trend in the rescue of repression by siREST, only 1 gene, *GANAB* (encoding for the neutral alpha-glucosidase AB in the glycan metabolism pathway) was significantly de-repressed (Figure S11). Overall, we confirmed that 3 out of 8 genes were bona fide REST target genes in hypoxia, from which so we could assume a similar percentage of the 201 newly identified REST target genes in hypoxia to be similarly genuine (38%, about 75 genes).

In order to test if the REST regulation of hypoxia repressed gene is direct or indirect, we searched for the consensus RE1 motif (Figure 5A) in the region from -2 kilobase pairs to +1 kilobase pairs from the transcription start site in the cohort of genes repressed by REST in hypoxia. In this search, we found high stringency consensus RE1 sites in just 12 genes (Figures 5B, C). This indicates that, in the majority of cases, REST is binding to distant or non-consensus sites. ChIP assays were performed to validate REST binding to the RE1 site on the promoters of *SYNJ1* (encoding for synaptojanin 1) and *GANAB*, two of the identified REST target genes in hypoxia (Figures 5B, D). For both *SYNJ1* and *GANAB*, we detected REST binding to their promoters in hypoxia (Figure 5D). For *SYNJ1*, we could observe a significant recruitment of REST in hypoxia over normoxia. We have also investigated whether the hypoxic REST-repressed genes from our RNA-seq are similar to published REST ChIP-Seq datasets from 4 different cell lines²⁷⁻²⁹. We found that, out of our list of 201 genes identified as REST-repressed in hypoxia, 36 were also REST target genes from the published ChIP-sequencing experiments (Figure 5E and Supplementary Table 4). This probably is still an under-estimation as there are no publically available REST ChIP-seq data performed on hypoxic cells.

Thus our unbiased approach using RNA-Seq revealed a previously unknown aspect of REST repression and identifies an extensive repertoire of genes regulated by REST in hypoxia (Figure 6), providing new insight into the mechanisms of gene repression in hypoxia.

Discussion

While HIF is considered to be the master regulator of increased gene expression in hypoxia²⁻⁴, the mechanisms of transcriptional repression in hypoxia are poorly understood¹¹. Our current study identifies REST as a key negative regulator of gene expression in hypoxia. We show it accumulates in the nucleus in response to hypoxia, and is responsible for the repression of approximately 20% of genes downregulated in hypoxia, thus acting as a counter-regulator to HIF-dependent gene expression.

REST is a phosphoprotein whose protein stability is tightly regulated during development. REST levels are reduced during neuronal development to allow the expression of REST target genes, while outside the nervous system, REST levels remain high to repress neuronal specific genes outside the nervous systems¹⁴. REST protein was stabilised in an *in vitro* model of ischemia (typically 10 to 30 minutes of glucose/oxygen deprivation followed by re-oxygenation, typically for 24 hours) due to an inhibition of its degradation³⁰. Our pulse-chase experiments with cycloheximide showed that hypoxia did not affect the stability of REST. Instead, there was increased nuclear localisation. **We also noticed that nuclear REST seemed to run at a smaller size. As REST is known to be phosphorylated and ubiquitylated²⁵, it is possible that post-translational modifications also control its translocation to the nucleus.**

Our experiments using REST siRNA in normoxic cells show that, in the basal state, only 164 genes of the detected normoxic transcriptome (23,284 genes) are repressed by REST (equivalent to approximately 0.7 % of the total transcriptome detected). Of the total of genes repressed in hypoxia (1013), we found that 201 gene are significantly repressed by REST in hypoxia (equivalent to approximately 20% of the total), **placing REST as a significant repressor of gene expression in hypoxia (Figure 6).** Using qRT-PCR, we had a 38% validation rate, suggesting about 75 of the 201 genes are expected to be *bona fide* REST targets in hypoxia. Furthermore, we found a small overlap (5 genes) between the REST-dependent genes in normoxia compared with the REST-dependent genes in hypoxia, indicating that REST regulates context-specific transcriptional networks. This has been previously described in other systems: for example REST was shown to regulate different target genes in neuronal stem cells versus embryonic stem cells³¹, embryonic stem cells vs epiblast stem cells³² and in response to ischemia in hippocampal neurons²¹. The factors leading to this differential recruitment are only starting to be uncovered, and may involve different co-factors, e.g. mSin3A³² and differences in the REST binding sequences^{31,33}. When we compared our normoxic/hypoxic REST target genes dataset with published ChIP-seq datasets, we found similarities of about 64% with the normoxic genes and 18% with the hypoxic genes. This

difference in proportions could be due to REST in hypoxia regulating more genes indirectly than by direct binding. This simple analysis reveals that there are layers of regulations in the control of gene expression and could open new research avenues into direct and indirect REST regulation.

Our ChIP and qRT-PCR experiments reveal two distinct binding patterns in 2 identified REST target genes *SYNJI* and *GANAB*. While REST was bound to both of their promoters in normoxia, we observed significant increased binding of REST to only the *SYNJI* promoter in hypoxia. We propose that, while REST recruitment is correlated to gene repression in hypoxia for some REST target genes (e.g. *SYNJI*), for other genes (e.g. *GANAB*) REST might be already bound to the target gene in a low affinity state, “poised to act,” with its repressive activity triggered only in hypoxia, possibly by co-factor recruitment. This is similar to the findings of Otto and colleagues ³⁴, which show that many of their identified REST target genes were still expressed while others were repressed, potentially due to their functional relevance to the cell. Interestingly *SYNJI* encodes for synaptotagmin, a protein involved in clathrin-mediated endocytosis. It could be speculated that this ATP-intensive process is marked for repression as soon as cellular levels of ATP become scarce.

Our RNA-seq experiment was performed on cells in hypoxia for 24 hours (to maximise gene repression ⁸) and we did not observe as dramatic an increase in HIF and its target genes as what we have previously reported in cells exposed to 8 hours of hypoxia ²³. Thus our experiments demonstrate the temporal nature of HIF signalling and its tight regulation by multiple transcription factors such as REST, NFκB ^{23,35} (which promotes *HIF1A* mRNA expression), or miR-155 ³⁶ (which de-stabilizes *HIF1A*). Of note, our previous study showed that NFκB is recruited to the *HIF1A* promoter with maximal binding at 8 hours in hypoxia and back to normoxic level after 24 hours ²³. It is thus possible that in the absence of REST (the repressor), we see increased *HIF1A* mRNA at 8 hours, but not at 24 hours, because one of the activators NFκB is less bound, or less active at this particular time point. The HIF signalling network is ripe for mathematical modelling to unravel its regulatory pathways ³⁷.

Gene ontology analysis suggests that REST plays an important role in the cellular adaptation to hypoxia by suppressing genes involved in proliferation (*DLX5*, *PRKCB* and *MET*), cell cycle progression (*TBX3*, *CABLES1*, *ARID3A* and *ARAP1*) and transcription, with transcription factors (TF) (*TCF12* and *LBX1*), regulators of TF activity (*MYCBP* and *ATF7IP*) and general transcription apparatus (*MMS19*, *CDK19* and *MED12*), consistent with our hypothesis that direct and indirect mechanisms contribute to REST gene repression in hypoxia. In addition REST also seems to

regulate important metabolic processes including the biosynthesis of lipids (*MBOAT2*, *GPAM*, *PIGN*, *AGPS* and *ACACA*) and nucleic acids (*PARP4*, *SKIV2L2*, *PLRG1* and *DHX37*). Interestingly REST also represses genes involved in protein catabolism (*FBXO18*, *NEDD4*, *ERLIN2* and *TRIP12*). Our own ChIP experiments confirmed the hypoxic regulation of *SYJN1* encoding synaptojanin 1, a protein involved in clathrin-mediated endocytosis. Thus repressing ATP-intensive processes in hypoxia is a plausible function for REST. Our lab has also shown a role for REST in the regulation of HIF-1 α , and thus could impact on HIF-regulated genes as well²³. Thus our transcriptomic analysis of hypoxic cells indicates that REST might play a key role in the regulation of proliferation and metabolic processes in hypoxia.

Some of the REST functions may be maladaptive in cancer. For example, REST-Dependent repression of cell adhesion genes in hypoxia (*ICAM5*, *ITGA8*, *SORBS1*, *LAMA4*, *LAMC1*, *LAMB1*, *COL14A1* and *COL4A5*) might be of importance for metastasis as hypoxic suppression of adhesion molecules in cancer cells has been proposed as a mechanism to allow hypoxic cancer cells to escape their stressed environment^{38,39}. Some of the previously described genes are part of cancer pathways (*RBI*, *MET*, *MYCBP*, *WNT5A*, *HDAC1*, *PRKCB*, *LAM4A*, *LAMC1* and *LAMB1*). The significance of the hypoxic repression of these genes by REST will be context- and cell-specific as REST may play both tumour suppressor and oncogenic roles, a feature it shares with NOTCH signalling⁴⁰, E-cadherin⁴¹ and MYC⁴². Indeed, a recent study⁴³ showed that knockdown of REST in normoxic prostate cancer cells can induce HIF signalling pathway, itself also implicated in cancer. In neuronal cancer, REST expression is high and has oncogenic properties by being anti-apoptotic and pro-tumorigenic^{15,44,45}. We speculate that the hypoxic microenvironment present in these tumours⁴⁶ could be a potential mechanism explaining REST over-expression. This is further supported by the observation that hypoxic neuroblastoma tumours and cells exposed to hypoxia (1% oxygen) down-regulate neuronal markers^{47,48}. REST is also implicated in stem cell renewal, and has been recently reported to accumulate in the nucleus of Marrow-isolated multilineage inducible (MIAMI) cells exposed to 3% oxygen⁴⁹.

Despite being widely regarded as an activator of gene expression in hypoxia, there are a few genes reported to be directly repressed by HIF-1 α in hypoxia (e.g. *CFTR*⁵⁰, *ADK*⁵¹ and *APC*⁵²). Other examples of hypoxia-induced transcriptional repressors include DEC1 (differentially expressed in chondrocytes protein 1) which represses MITF (Microphthalmia-associated transcription factor)⁵³ and Bach1 (BTB and CNC homology 1, basic leucine zipper transcription factor 1) which represses HO-1 (heme oxygenase-1)⁵⁴. However, none of these appear to regulate as broad a range of genes in hypoxia as REST. Indeed, our data indicates that REST regulates 20% of genes

repressed in hypoxia, which may be a key and previously unappreciated feature of the hypoxic response. REST-repressed genes regulate biosynthetic metabolism, cell cycle and proliferation indicating a major role in the cellular adaptation to hypoxia. The whole set of 201 genes repressed in hypoxia holds exciting possibilities regarding the discovery of novel mechanisms in the hypoxic response mediated by transcriptional repression. In summary, our findings have identified REST as a key regulator underlying gene repression and cellular adaptation in hypoxia.

Methods

Cell culture

Human embryonic kidney cells (HEK-293) were cultured in Dulbecco's modified eagle medium (DMEM, high glucose 4.5 g/L without pyruvate) supplemented with 10% foetal calf serum (FCS) and 100 U/mL penicillin-streptomycin (PS). Human non-tumorigenic mammary epithelial cells (MCF10A) were grown in DMEM/F12 with 5% horse serum, 100 U/mL PS, 2 mM L-Glutamine, 10µg/mL insulin, 20ng/mL Epidermal Growth Factor (EGF), 0,5µg/mL hydrocortisone and 100ng/ml cholera toxin. All cells were obtained from the American Type Culture Collection (ATCC). All reagents for cell culture were from Gibco (Life Technologies, Calrsbad, CA, USA), unless otherwise stated. Cells were exposed to hypoxia using pre-equilibrated media and maintained in standard normobaric hypoxic conditions (1% O₂, 5% CO₂ and 94% N₂) in a hypoxia chamber (Coy Laboratories, Grass Lake, Michigan, USA). Normoxic controls were maintained at atmospheric O₂ levels (21% O₂, 5% CO₂ and 74% N₂) in a tissue culture incubator. Re-oxygenation was performed by bringing hypoxic cells to the tissue culture incubator for the indicated time points.

Cloning

All reagents were from New England Biolabs (NEB) unless otherwise stated. The pRE1-TK-GLuc REST responsive construct was developed by incorporating the RE1 of the *SCN2A* gene⁵⁵ into the vector pTK-GLuc (NEB, N8084S). Primers were designed to amplify the RE1 using primerBLAST (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>). At the 5'-end of each primer additional nucleotides were introduced containing the restriction sites (capital letters) and additional nucleotides to facilitate digestion by the restriction endonucleases, a NotI restriction site 5'-tattGCGGCCGC-3' was introduced into the forward primer 5'-TTTCTCTATCGATAGGTACAGGCA -3' and a XhoI restriction site 5'-ctattCTCGAG-3' was introduced into the reverse primer 5'- GTAATTCCAATTGTGACCAGGA-3'. PCR-cloning, digestion, ligation and sub-cloning were performed using standard molecular biology protocols. The pHRE-MP-GLuc HIF responsive construct has been previously described⁵⁶. The pCMV-CLuc construct was from NEB (N0321S, pCMV-CLuc 2). Plasmid sequencing was performed by MWG Eurofins, Germany.

Gaussia luciferase assay, transient and stable transfections

Gaussia luciferase assays were performed as previously described⁵⁶. Briefly, at the selected time points, 10 µL of media was collected from the supernatant and stored at -20°C. Gaussia luciferase activity was measured using the Biolum Gaussia luciferase Flex Assay kit (NEB) in a plate reader

(Synergy HT, Biotek) and normalized to the luciferase activity of the secreted cypridina luciferase under the control of a constitutive CMV promoter (pCMV-CLuc) or protein concentration. Details and validation of the constructs are shown in Figure S4.

Cell transfection with siRNA

Transient transfection with siControl (sc-37007, SCB) and siREST (s11932, Life Technologies) were performed as previously described⁵⁷. Transfections with siRNA to be used in luciferase assays were performed in 24 well plates, as described above. All other experiments were performed on 6 well plates unless otherwise stated. In a typical experiment 200 K cells were seeded on 6 well plates and allowed to grow until approximately 60 % confluent, at this time cells were transfected with 2 μ L of Lipo, 100 μ L Optimem and a pre-optimized amount of overexpressing construct (100 ng) or siRNA (100 pmol). **On the day after transfection, cells were exposed to hypoxic or normoxic media. Transfection time with siRNA was kept constant for all experimental conditions. For RNA extraction experiments, siRNA was incubated for 48 hrs. For the preparation of whole cell protein extracts, siRNA was incubated for 72 hrs.**

Cycloheximide (CHX) pulse chase experiments

For the experiments where CHX-pulse chase was used to determine the protein stability of HIF-1 α , REST and β -Actin in normoxia and hypoxia, 750,000 HEK293 cells were seeded on 6 cm dishes. The following day cells were conditioned to 24 hours of hypoxia or normoxia, after which cells were subjected to a pulse-chase treatment of cycloheximide (5 μ g/mL) to block translation, and address the stability of the proteins in normoxia and hypoxia. Time points of CHX treatment after normoxic or hypoxic incubation included 0, 1, 2, 4, 8, 24 and 48 hours.

qRT-PCR

cDNA was synthesized from 1 μ g of RNA using MMLV (Promega), and amplified using the Prism 7900HT sequence detection system (Applied Biosystems, Foster City, CA) under default conditions. The mRNA relative expression was calculated by the $\Delta\Delta$ Ct method by normalizing the Ct of the samples to that of 18S rRNA (TaqMan Universal PCR Master Mix with the primer 18S rRNA-Euka, 4310893E, Life Technologies), followed by normalization to the control condition. The following REST qRT-PCR primers were used:

F: CGCCAGAGGGTGAAACTTTA; R: ATCCACAGCCATGAAGGAAG

Western blot

All reagents were from Sigma unless otherwise stated. Standard protocols were used as previously described^{57,58}. Mouse anti- β -actin, 1:10000, Sigma, A5441; Mouse anti-HIF-1 α , 1:1000, BD Pharmingen, 610958; Rabbit anti-REST, 1:1000, Abcam, ab28018; Rabbit anti-Lamin A/C, 1:1000, Cell Signalling, 2032; Mouse anti- α -Tubulin, 1:2000, SCB, sc-8035.

ChIP assays

ChIP assays were performed as previously described⁵⁹.

The following ChIP qRT-PCR primers were used:

SYNJ1, F: TCCAGACACTCAGACTAGGAACTC, R: CCTGAAGAGCTGTCCATGGT, Probe: CCGTTTGCTGGGCTGTCGAC.

The following antibodies were used:

Rabbit REST, 2 μ g, Millipore, Rabbit IgG, 5 μ g, Millipore, PP64B.

Cell death using flow cytometry (YO-PRO and PI stains) and trypan blue exclusion assays.

Trypan Blue (Sigma), was added 1:1 V/V to the cells, and live/dead cells were counted in a haemocytometer, by counting the Trypan blue permeabilised cells (dead), and the bright cells (alive), that were not Trypan Blue permeabilised. Data is represented as a percentage of the total population. Flow Cytometry staining to detect live and dead cells (early, late and end stage apoptotic cells) was performed using standard protocols. Briefly, cells were detached with Accutase solution, and stained with PI/YO-PRO, and counted in an Accuri C6 cytometer. Results are shown as the percentage of the total population under each condition.

REST mRNA expression profiling

Human REST mRNA expression level according to anatomical distribution, or in different cell lines, was compiled from publically available microarray datasets using the Genevestigator software (<https://genevestigator.com/gv/>).

RNA sequencing

HEK293 (100 thousand cells) were seeded on 6 well plates, transfected the next day with REST-RNAi or Ctrl-RNAi in 1 μ L of Lipofectamine 2000. In the day after transfection cells were exposed to 1% oxygen using pre-conditioned media for 24 hours. RNA was isolated using QIAgen RNeasy columns as per supplier instructions. Total RNA electropherograms were used to evaluate RNA integrity, with integrity number for all samples being of the highest quality, RIN=10, before being sent to the sequencing facility. Library construction began with 1 μ g of total RNA. Since the protocol utilized was based on polyA capture, RNA was visualized on a BioRad Experion to

insure RNA Quality Index values were greater than 8. For library construction the Illumina TruSeq v2 mRNA kit was utilized. This included also the poly(A)+ RNA isolation and DNase treatment. The protocol was followed according to manufacturer's instructions except the final number of PCR cycles was 12 and not 15. Fifty base pair single end reads were prepared. Following construction, libraries were visualized by Bioanalyzer (Agilent) using the High Sensitivity Chip and quantified for pooling and sequencing using Kapa Biosystems qPCR quantitation kit according to manufacturer's instructions. For sequencing libraries were diluted to 16 pM then applied to a V3 flowcell using the Illumina cBot according to manufacturer's instructions. Sequencing was carried out on the Hi-Seq 2000 using HSCS v 1.5.15.1. Image analysis and base calling were carried out using RTA 1.13.48.0, and deconvolution and fastq file production were performed with CASAVA 1.8.2. Quality control metrics were obtained with Picard after Illumina FASTQ files alignment to the human genome hg19 reference sequence (GRCh37) using TopHat v2.0.8b ⁶⁰. Raw sequencing files and derived alignment files data are available in the ArrayExpress database (<https://www.ebi.ac.uk/arrayexpress/>) under the accession number E-MTAB-2580.

RNA sequencing bioinformatics analysis

Illumina FASTQ files were aligned to the human genome hg19 reference sequence (GRCh37) using TopHat v2.0.8b ⁶⁰ (including Bowtie 2.1.0.0 and Samtools 0.1.19.0) with the -no-coverage-search option and the -G option to use a transcriptome index computed from the UCSC hg19 genes.gtf annotation. Transcript abundance and differential expression was estimated by direct comparison of normalized transcript counts from pairs of experimental conditions using the Cufflinks Cuffdiff2 program (v2.1.1) ⁶¹ with default options and the UCSC hg19 genes.gtf annotation. To control for false positives, a second estimate of gene expression was computed using the HTSeq-count script (v0.6.1; ⁶² followed by differential expression analysis with the Bioconductor edgeR package (v3.6.1) ⁶³, which uses a generalized linear model likelihood ratio test. Benjamini and Hochberg's algorithm is used to control the false discovery rate (FDR) due to multiple testing ⁶⁴, those genes with FDR (q-value) < 0.05 were considered differentially expressed. The February 2009 human reference sequence (GRCh37) was produced by the Genome Reference Consortium (<http://www.ncbi.nlm.nih.gov/projects/genome/assembly/grc/>).

Analysis of the promoter of REST-Dependent hypoxia repressed genes

Genomic sequences of genes identified as differentially regulated ("REST-Dependent genes repressed in hypoxia") were extracted from the USCS Genome Browser database, selecting 2000 bases upstream (5') of the transcriptional start site (TSS) and 1000 bases 3' of the TSS. When

multiple TSS sites were annotated for a single gene, the 3Kb region was selected for all of them. The motif for the RE1 site from the JASPAR database (MA0138.2) was used as a frequency matrix and compared to the 3Kb sequences by the pssm.search method using the Biopython Bio.motifs module, with a pseudocount of 0.5 and Normal background (equal frequency of G:A:T:C bases). The log odds threshold was set to 15, which would yield an expected false positive rate of approximately 0.05 against random background sequences. The sensitivity of this method was calibrated against equal sized sets of 3Kb random sequences (using RSA-Tools) and against the target sequences with the RE1 motif bases shuffled (using EMBOSS).

We found 12 high stringency RE1 motifs (log odds score of 15) in the promoter region (-2Kb/+1Kb) of REST-dependent genes repressed in hypoxia (Figure 6B). This number of hits is much higher than what could be found in control experiments, when the RE1 motif was searched against an equal sized set of random sequences (2.4 ± 0.54 gene hits, Figure 6C) or randomly shuffling the promoter sequences from the REST genes (0.4 ± 0.54 gene hits, Figure 6C). This analysis strengthens the point that the REST-dependent genes repressed in hypoxia have consensus RE1 motifs in the -2Kb/+1Kb region of their promoter in a greater frequency than one would expect by chance.

Functional analysis of gene datasets using PANTHER and DAVID

PANTHER was used for the functional annotation of gene lists (Figures S1A-J) and for the statistical overrepresentation tests (Figures 1C-H), significance threshold was set as $p < 0.05$ for the binomial test^{65,66}. This is a hypergeometric test which first assigns each gene to one or more GO or PANTHER categories, and then tests if there are more genes in a particular category than those one would expect by random chance in a dataset of similar size. DAVID Functional Annotation Clustering was performed with default settings, the significance threshold of this gene-enrichment analysis was set as $p < 0.05$ for the EASE Score (a modified Fisher Exact p-Value)^{67,68}.

Statistical analysis

All experiments were performed at least 3 independent times. Data is shown as mean \pm SEM. Statistical significance was tested in Prism (Graphpad), using Student's t test for the comparison of two data sets or ANOVA for more than two datasets. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$.

Author contributions

M.A.S.C., C.T.T. and A.C. designed the study and experiments, M.A.S.C., M.M., B.C., M.C.M., M.M.T., A.C.S., C.E.K., S.B., A.W., D.C., A.W., Z.F. and C.C.S. performed experiments, M.A.S.C., M.M., B.C., S.B., C.T.T., L.M.A.R., K.A.N., C.G., K.J.M., E.P.C. and A.C. analysed the data, M.A.S.C., C.T.T, S.B. and A.C. wrote the manuscript.

Competing financial interests

The authors declare no competing interest.

Acknowledgements

This work was supported by the Science Foundation Ireland under Grant No. 06/CE/B1129. We would like to thank Professor Louis B. Hersh (University of Kentucky, USA) for the kind gift of the msREST-FLAG and Dr Ian Wood (University of Leeds, UK) for the rat *SCN2A* RE1 sequence.

References

- 1 Semenza, G. L. Hypoxia-inducible factors in physiology and medicine. *Cell* **148**, 399-408, doi:10.1016/j.cell.2012.01.021 (2012).
- 2 Elvidge, G. P. *et al.* Concordant regulation of gene expression by hypoxia and 2-oxoglutarate-dependent dioxygenase inhibition - The role of HIF-1 alpha, HIF-2 alpha, and other pathways. *J Biol Chem* **281**, doi:10.1074/jbc.M511408200 (2006).
- 3 Manalo, D. J. *et al.* Transcriptional regulation of vascular endothelial cell responses to hypoxia by HIF-1. *Blood* **105**, 659-669, doi:10.1182/blood-2004-07-2958 (2005).
- 4 Benita, Y. *et al.* An integrative genomics approach identifies Hypoxia Inducible Factor-1 (HIF-1)-target genes that form the core response to hypoxia. *Nucleic Acids Res* **37**, 4587-4602, doi:10.1093/nar/gkp425 (2009).
- 5 Schofield, C. J. & Ratcliffe, P. J. Oxygen sensing by HIF hydroxylases. *Nat Rev Mol Cell Biol* **5**, 343-354, doi:10.1038/nrm1366 (2004).
- 6 Prabhakar, N. R. & Semenza, G. L. Adaptive and maladaptive cardiorespiratory responses to continuous and intermittent hypoxia mediated by hypoxia-inducible factors 1 and 2. *Physiol Rev* **92**, doi:10.1152/physrev.00030.2011 (2012).
- 7 Semenza, G. L. Defining the role of hypoxia-inducible factor 1 in cancer biology and therapeutics. *Oncogene* **29**, doi:10.1038/onc.2009.441 (2010).
- 8 Starmans, M. H. W. *et al.* The prognostic value of temporal in vitro and in vivo derived hypoxia gene-expression signatures in breast cancer. *Radiotherapy and Oncology* **102**, 436-443, doi:10.1016/j.radonc.2012.02.002 (2012).
- 9 Cummins, E. P. & Taylor, C. T. Hypoxia-responsive transcription factors. *Pflugers Archiv-European Journal of Physiology* **450**, 363-371, doi:10.1007/s00424-005-1413-7 (2005).
- 10 Schoedel, J. *et al.* High-resolution genome-wide mapping of HIF-binding sites by ChIP-seq. *Blood* **117**, E207-E217, doi:10.1182/blood-2010-10-314427 (2011).
- 11 Chi, J. T. *et al.* Gene expression programs in response to hypoxia: Cell type specificity and prognostic significance in human cancers. *Plos Medicine* **3**, 395-409, doi:10.1371/journal.pmed.0030047 (2006).
- 12 Gaston, K. & Jayaraman, P. S. Transcriptional repression in eukaryotes: repressors and repression mechanisms. *Cellular and Molecular Life Sciences* **60**, 721-741, doi:10.1007/s00018-003-2260-3 (2003).
- 13 Tadepally, H. D., Burger, G. & Aubry, M. Evolution of C2H2-zinc finger genes and subfamilies in mammals: Species-specific duplication and loss of clusters, genes and effector domains. *BMC Evol Biol* **8**, doi:10.1186/1471-2148-8-176 (2008).
- 14 Ooi, L. & Wood, I. C. Chromatin crosstalk in development and disease: lessons from REST. *Nat Rev Genet* **8**, doi:10.1038/nrg2100 (2007).
- 15 Negrini, S., Prada, I., D'Alessandro, R. & Meldolesi, J. REST: an oncogene or a tumor suppressor? *Trends Cell Biol* **23**, 289-295, doi:10.1016/j.tcb.2013.01.006 (2013).
- 16 Schoenherr, C. J., Paquette, A. J. & Anderson, D. J. Identification of potential target genes for the neuron-restrictive silencer factor. *Proceedings of the National Academy of Sciences of the United States of America* **93**, 9881-9886, doi:10.1073/pnas.93.18.9881 (1996).
- 17 Kuwahara, K. *et al.* The neuron-restrictive silencer element-neuron-restrictive silencer factor system regulates basal and endothelin 1-inducible atrial natriuretic peptide gene expression in ventricular myocytes. *Molecular and Cellular Biology* **21**, 2085-2097, doi:10.1128/mcb.21.6.2085-2097.2001 (2001).

- 18 Cheong, A. *et al.* Downregulated REST transcription factor is a switch enabling critical potassium channel expression and cell proliferation. *Mol Cell* **20**, 45-52, doi:10.1016/j.molcel.2005.08.030 (2005).
- 19 Scholl, T., Stevens, M. B., Mahanta, S. & Strominger, J. L. A zinc finger protein that represses transcription of the human MHC class II gene, DPA(1,2). *Journal of Immunology* **156**, 1448-1457 (1996).
- 20 Lu, T. *et al.* REST and stress resistance in ageing and Alzheimer's disease. *Nature* **507**, 448-454, doi:10.1038/nature13163 (2014).
- 21 Noh, K.-M. *et al.* Repressor element-1 silencing transcription factor (REST)-dependent epigenetic remodeling is critical to ischemia-induced neuronal death. *Proceedings of the National Academy of Sciences of the United States of America* **109**, E962-971, doi:10.1073/pnas.1121568109 (2012).
- 22 Calderone, A. *et al.* Ischemic insults derepress the gene silencer REST in neurons destined to die. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **23**, 2112-2121 (2003).
- 23 Cavadas, M. A. *et al.* REST mediates resolution of HIF-dependent gene expression in prolonged hypoxia. *Scientific reports* **5**, 17851, doi:10.1038/srep17851 (2015).
- 24 Liu, Z., Liu, M., Niu, G., Cheng, Y. & Fei, J. Genome-wide identification of target genes repressed by the zinc finger transcription factor REST/NRSF in the HEK 293 cell line. *Acta Biochim Biophys Sinica* **41**, 1008-1017, doi:10.1093/abbs/gmp095 (2009).
- 25 Westbrook, T. F. *et al.* SCF beta-TRCP controls oncogenic transformation and neural differentiation through REST degradation. *Nature* **452**, 370-U311, doi:10.1038/nature06780 (2008).
- 26 Wagoner, M. P. *et al.* The Transcription Factor REST Is Lost in Aggressive Breast Cancer. *Plos Genetics* **6**, doi:10.1371/journal.pgen.1000979 (2010).
- 27 Johnson, D. S., Mortazavi, A., Myers, R. M. & Wold, B. Genome-wide mapping of in vivo protein-DNA interactions. *Science* **316**, 1497-1502, doi:10.1126/science.1141319 (2007).
- 28 Johnson, R. *et al.* REST regulates distinct transcriptional networks in embryonic and neural stem cells. *PLoS biology* **6**, e256, doi:10.1371/journal.pbio.0060256 (2008).
- 29 Satoh, J., Kawana, N. & Yamamoto, Y. ChIP-Seq Data Mining: Remarkable Differences in NRSF/REST Target Genes between Human ESC and ESC-Derived Neurons. *Bioinformatics and biology insights* **7**, 357-368, doi:10.4137/BBI.S13279 (2013).
- 30 Kaneko, N., Hwang, J.-Y., Gertner, M., Pontarelli, F. & Zukin, R. S. Casein Kinase 1 Suppresses Activation of REST in Insulted Hippocampal Neurons and Halts Ischemia-Induced Neuronal Death. *Journal of Neuroscience* **34**, 6030-6039, doi:10.1523/jneurosci.4045-13.2014 (2014).
- 31 Johnson, R. *et al.* REST Regulates Distinct Transcriptional Networks in Embryonic and Neural Stem Cells. *Plos Biology* **6**, 2205-2219, doi:10.1371/journal.pbio.0060256 (2008).
- 32 Seki, M. *et al.* A Comparison of the Rest Complex Binding Patterns in Embryonic Stem Cells and Epiblast Stem Cells. *Plos One* **9**, doi:10.1371/journal.pone.0095374 (2014).
- 33 Rockowitz, S. *et al.* Comparison of REST Cistromes across Human Cell Types Reveals Common and Context-Specific Functions. *Plos Computational Biology* **10**, doi:10.1371/journal.pcbi.1003671 (2014).
- 34 Otto, S. J. *et al.* A new binding motif for the transcriptional repressor REST uncovers large gene networks devoted to neuronal functions. *The Journal of neuroscience : the*

- official journal of the Society for Neuroscience **27**, 6729-6739, doi:10.1523/JNEUROSCI.0091-07.2007 (2007).
- 35 BelAiba, R. S. *et al.* Hypoxia up-regulates hypoxia-inducible factor-1 alpha transcription by involving phosphatidylinositol 3-kinase and nuclear factor kappa B in pulmonary artery smooth muscle cells. *Mol Biol Cell* **18**, 4691-4697, doi:10.1091/mbc.E07-04-0391 (2007).
 - 36 Bruning, U. *et al.* MicroRNA-155 Promotes Resolution of Hypoxia-Inducible Factor 1 alpha Activity during Prolonged Hypoxia. *Mol Cell Biol* **31**, doi:10.1128/mcb.01276-10 (2011).
 - 37 Cavadas, M. A. S., Nguyen, L. K. & Cheong, A. Hypoxia-inducible factor (HIF) network: insights from mathematical models. *Cell Commun Signal* **11**, doi:10.1186/1478-811X-11-42 (2013).
 - 38 Hasan, N. M., Adams, G. E., Joiner, M. C., Marshall, J. F. & Hart, I. R. Hypoxia facilitates tumour cell detachment by reducing expression of surface adhesion molecules and adhesion to extracellular matrices without loss of cell viability. *British Journal of Cancer* **77**, 1799-1805, doi:10.1038/bjc.1998.299 (1998).
 - 39 Pouyssegur, J., Dayan, F. & Mazure, N. M. Hypoxia signalling in cancer and approaches to enforce tumour regression. *Nature* **441**, 437-443, doi:10.1038/nature04871 (2006).
 - 40 Lobry, C., Oh, P. & Aifantis, I. Oncogenic and tumor suppressor functions of Notch in cancer: it's NOTCH what you think. *Journal of Experimental Medicine* **208**, 1931-1935, doi:10.1084/jem.20111855 (2011).
 - 41 Rodriguez, F. J., Lewis-Tuffin, L. J. & Anastasiadis, P. Z. E-cadherin's dark side: Possible role in tumor progression. *Biochimica Et Biophysica Acta-Reviews on Cancer* **1826**, 23-31, doi:10.1016/j.bbcan.2012.03.002 (2012).
 - 42 Uribealago, I., Aznar Benitah, S. & Di Croce, L. From oncogene to tumor suppressor The dual role of Myc in leukemia. *Cell Cycle* **11**, 1757-1764, doi:10.4161/cc.19883 (2012).
 - 43 Lin, T. P. *et al.* REST reduction is essential for hypoxia-induced neuroendocrine differentiation of prostate cancer cells by activating autophagy signaling. *Oncotarget*, doi:10.18632/oncotarget.8433 (2016).
 - 44 Conti, L. *et al.* REST Controls Self-Renewal and Tumorigenic Competence of Human Glioblastoma Cells. *Plos One* **7**, doi:10.1371/journal.pone.0038486 (2012).
 - 45 Huang, Z. & Bao, S. Ubiquitination and deubiquitination of REST and its roles in cancers. *Febs Letters* **586**, doi:10.1016/j.febslet.2012.04.052 (2012).
 - 46 Kaur, B. *et al.* Hypoxia and the hypoxia-inducible-factor pathway in glioma growth and angiogenesis. *Neuro-Oncology* **7**, 134-153, doi:10.1215/s1152851704001115 (2005).
 - 47 Lofstedt, T. *et al.* Induction of ID2 expression by hypoxia-inducible factor-1 - A role in dedifferentiation of hypoxic neuroblastoma cells. *Journal of Biological Chemistry* **279**, 39223-39231, doi:10.1074/jbc.M402904200 (2004).
 - 48 Jogi, A. *et al.* Hypoxia alters gene expression in human neuroblastoma cells toward an immature and neural crest-like phenotype. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 7021-7026, doi:10.1073/pnas.102660199 (2002).
 - 49 Rios, C. *et al.* Low Oxygen Modulates Multiple Signaling Pathways, Increasing Self-Renewal, While Decreasing Differentiation, Senescence, and Apoptosis in Stromal MIAMI Cells. *Stem cells and development* **25**, 848-860, doi:10.1089/scd.2015.0362 (2016).

- 50 Zheng, W. *et al.* Hypoxia inducible factor-1 (HIF-1)-mediated repression of cystic fibrosis transmembrane conductance regulator (CFTR) in the intestinal epithelium. *Faseb Journal* **23**, 204-213, doi:10.1096/fj.08-110221 (2009).
- 51 Morote-Garcia, J. C., Rosenberger, P., Kuhlicke, J. & Eltzschig, H. K. HIF-1-dependent repression of adenosine kinase attenuates hypoxia-induced vascular leak. *Blood* **111**, 5571-5580, doi:10.1182/blood-2007-11-126763 (2008).
- 52 Newton, I. P., Kenneth, N. S., Appleton, P. L., Naethke, I. & Rocha, S. Adenomatous Polyposis Coli and Hypoxia-inducible Factor-1 alpha Have an Antagonistic Connection. *Molecular Biology of the Cell* **21**, 3630-3638, doi:10.1091/mbc.E10-04-0312 (2010).
- 53 Feige, E. *et al.* Hypoxia-induced transcriptional repression of the melanoma-associated oncogene MITF. *Proceedings of the National Academy of Sciences of the United States of America* **108**, E924-E933, doi:10.1073/pnas.1106351108 (2011).
- 54 Kitamuro, T. *et al.* Bach1 functions as a hypoxia-inducible repressor for the heme oxygenase-1 gene in human cells. *Journal of Biological Chemistry* **278**, 9125-9133, doi:10.1074/jbc.M209939200 (2003).
- 55 Mori, N., Schoenherr, C., Vandenberg, D. J. & Anderson, D. J. A common silencer element in the SCG10 and type II Na⁺ channel genes binds a factor present in nonneuronal cells but not in neuronal cells. *Neuron* **9**, 45-54, doi:10.1016/0896-6273(92)90219-4 (1992).
- 56 Cavadas, M. A. S. & Cheong, A. Monitoring of Transcriptional Dynamics of HIF and NF kappa B Activities. *Bioluminescent Imaging: Methods and Protocols* **1098**, 97-105, doi:10.1007/978-1-62703-718-1_8 (2014).
- 57 Nguyen, L. K. *et al.* A dynamic model of the hypoxia-inducible factor 1 α (HIF-1 α) network. *J Cell Sci* **126**, 1454-1463, doi:10.1242/jcs.119974 (2013).
- 58 Scholz, C. C. *et al.* Regulation of IL-1 beta-induced NF-kappa B by hydroxylases links key hypoxic and inflammatory signaling pathways. *Proc Natl Acad Sci USA* **110**, 18490-18495, doi:10.1073/pnas.1309718110 (2013).
- 59 Nguyen, L. K., Cavadas, M. A. S., Kholodenko, B. N., Frank, T. D. & Cheong, A. Species differential regulation of COX2 can be described by an NF kappa B-dependent logic AND gate. *Cell Mol Life Sci* **72**, 2431-2443, doi:10.1007/s00018-015-1850-1 (2015).
- 60 Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biology* **14**, doi:10.1186/gb-2013-14-4-r36 (2013).
- 61 Trapnell, C. *et al.* Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nature Biotechnology* **31**, 46-+, doi:10.1038/nbt.2450 (2013).
- 62 Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166-169, doi:10.1093/bioinformatics/btu638 (2015).
- 63 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140, doi:10.1093/bioinformatics/btp616 (2010).
- 64 Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Methodological* **57**, 289-300 (1995).
- 65 Mi, H., Muruganujan, A. & Thomas, P. D. PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic Acids Research* **41**, D377-D386, doi:10.1093/nar/gks1118 (2013).

- 66 Mi, H., Muruganujan, A., Casagrande, J. T. & Thomas, P. D. Large-scale gene
function analysis with the PANTHER classification system. *Nature Protocols* **8**,
1551-1566, doi:10.1038/nprot.2013.092 (2013).
- 67 Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources. *Nature Protocols* **4**, 44-57,
doi:10.1038/nprot.2008.211 (2009).
- 68 Huang, D. W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools:
paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids
Research* **37**, 1-13, doi:10.1093/nar/gkn923 (2009).

Figure legends

Figure 1. Hypoxia induced gene expression changes assessed by genome wide sequencing. (A) Volcano plot showing changes in gene expression due to hypoxia (1% O₂) plotted against significance. Each dot represents the mean fold change for a single gene with induced genes as green dots, repressed as red and unchanged as black. Horizontal blue dashed line indicates $q < 0.05$. Cells were treated with Ctrl_RNAi before hypoxia exposure and RNA-seq. (B) Mean fold change values for well described hypoxia induced (green bars) and repressed genes (red bars). (C, D) Enriched GO Biological processes.

Figure 2. REST is a hypoxia sensitive transcription factor. (A) REST mRNA expression compiled from publically available microarray datasets of different tissues and cell lines. The graph was compiled with Genevestigator and shows the human REST mRNA expression levels (xx axis) and number of samples (yy). HEK293 cells were exposed to hypoxia (1% O₂) for the indicated time points, nuclear extracts (B), cytoplasmic (D) and whole cell extracts (F) were prepared and immunoblotted as indicated for REST, HIF-1 α , Lamin A/C (nuclear marker) and α -Tubulin (cytoplasmic marker) (n = 4-5). (C, E, G) Densitometric analysis of B, D and F, respectively. (H) HEK293 cells where exposed to hypoxia (1% O₂) for 16 hours followed by re-oxygenation (21% O₂) for 1 and 16 hours, cytoplasmic and nuclear extracts where prepared and immunoblotted as indicated (N=4). (I) Cells were transfected with a vector expressing *Gaussia* luciferase under the control of the TK promoter (pTK-GLuc) for constitutive expression, or with the REST responsive pRE1-TK-GLuc construct. REST repressor activity = $RLU_{pTK-GLuc} / RLU_{pRE1-TK-GLuc}$ (N=3). (J) HEK293 cells were exposed to hypoxia (1% O₂) for the indicated time points, mRNA was collected and analysed by qRT-PCR, n = 6-10. Data are represented as mean \pm SEM. * = $p < 0.05$, ** = $p < 0.01$, significant fold change over 21% O₂.

Figure 3. REST regulates hypoxia repressed genes. (A) Overview of the RNA-seq approach used to reveal the contribution of REST to gene repression in hypoxia. (B) REST mRNA expression from the 3 independent RNA-seq experiments after treatment with hypoxia or REST-RNAi treatment. (C) REST-RNAi treatment on cells exposed to normoxia (21% oxygen) for 24 hours led to a significant increase in the expression of well-characterised REST target genes (shown as Cuffdiff mean fold changes (log base 2) of 3 independent experiments). (D) RNA-Seq analysis revealed 201 REST-Dependent genes

(Red) among the repressed genes (Blue). Data shown as a Volcano plot showing changes in gene expression due to hypoxia (1% O₂) plotted against significance. Only genes with statistically significant fold changes are shown. Each dot represents the mean fold change for a single gene with induced-genes as green dots (“Hypoxia induced”), repressed-genes as blue (“Hypoxia repressed”), and the genes that were not repressed by hypoxia in the presence of siRNA against REST shown as red dots (“REST-Dependent”). **(E)** Heat-map showing genes repressed by hypoxia (left column, Ctrl-RNAi), that were not repressed upon REST-RNAi treatment (right column, REST-RNAi). Data is presented as fold change to hypoxia of the top 20 genes most repressed by hypoxia among the REST-dependent genes. **(F)** Summary of the gene clusters found to be uniquely repressed by REST-Independent mechanisms (Blue), REST-Dependent mechanisms (Red) and the common clusters between the two mechanisms (Grey). Clusters were obtained using DAVID Functional Annotation Clustering together with manual curation of the results. For a detailed description of the annotation terms and gene groups associated with each cluster see Table S2.

Figure 4. REST regulates different transcriptional networks in normoxia and hypoxia.

(A) There is limited overlap (5 genes) between REST-Dependent genes repressed in normoxia (164, dark circle) and REST-Dependent genes repressed in hypoxia (201, red circle). Circular diagrams show the distribution of enriched GO biological processes by PANTHER overrepresentation test. Significance was set at $p < 0.05$. The top 20 genes by fold-change upon REST-RNAi treatment for the REST-Dependent genes repressed in normoxia **(B)** and REST-dependent genes repressed in hypoxia **(C)** are shown. **(D, E)** qRT-PCR was used to validate the mRNA expression of the REST target genes *RAB3C* and *CYP11B1* in cells exposed for 24 hours to normoxia (21% oxygen) or hypoxia (1% oxygen), in the presence or absence of an siRNA targeting REST. **(F)** Of the 851 genes induced in hypoxia (blue circle), 29 were further induced by REST siRNA treatment. Circular diagrams show the distribution of enriched GO biological processes by PANTHER overrepresentation test.


Figure 5. REST represses genes in hypoxia by direct binding to an RE1 site on their promoter regions.

(A) Consensus RE1 binding site (JASPAR matrix representation). **(B)** Identification of 12 high stringency RE1 sites in the proximal promoter (-2000 bp to +1000 bp kilobase) of REST-dependent hypoxia repressed genes. **(C)** The number of RE1 sites identified by our informatics analysis is compared to the number of RE1 sites that could be expected by random chance (see Materials and Methods for details. Data is presented as


mean \pm SD. Error bars represent the SD from the 500 simulations. **(D)** ChIP assay for REST binding in normoxia and hypoxia (24 hours) to the RE1 found in the proximal promoter of the *GANAB* and *SYNJ1* gene. **(E)** Overlap between the number of genes identified to be REST target genes in normoxia or hypoxia, with the REST target genes identified by ChIP-seq in published datasets. Data is represented as mean \pm SEM. * = $p < 0.05$, significant over normoxic control and IgG. *n.d.* = none detected, for some of the biological replicates.

Figure 6. REST is a master regulator of gene repression in hypoxia and regulates different transcriptional networks depending on oxygen availability. Our analysis showed that REST repressed 20% of the hypoxia-repressed genes, and only 0.7% of all genes repressed in normoxia. Furthermore there is only a limited overlap between the 2 transcriptional networks (5 genes), and most of the genes repressed by REST in normoxia are still repressed by REST under hypoxia. Thus, rather than a shift on the transcriptional repertoire of REST target genes from normoxia to hypoxia, we show that there is a recruitment to new target genes in hypoxia.


A




B




C




D




A




H




B




C




D




E




I




F




G




J




A




B




C




D




E



F





A



neurological system process
ion transport
cell communication
Other Biological processes

REST-
repressed in
normoxia

REST-
repressed in
hypoxia



metabolic process
primary metabolic process
cell adhesion
biological adhesion

B


REST-repressed genes in normoxia (21 % oxygen)			
Gene symbol	Gene name	siREST-Fold change (log base 2)	p-Value
CPLX2	complexin 2	6,7	9,7E-205
CHGA	chromogranin A	5,3	2,9E-176
UNC13A	unc-13 homolog A (C. elegans)	4,6	1,5E-166
SCG3	secretogranin III	4,7	1,2E-139
CHGB	chromogranin B	4,9	3,8E-90
CPLX1	complexin 1	2,6	1,0E-87
ACTL6B	actin-like 6B	5,0	3,4E-82
AP3B2	adaptor-related protein complex 3, beta 2 subunit	2,6	1,7E-77
XKR7	XK, Kell blood group complex subunit-related family, member 7	6,7	8,6E-62
LOC254559	hypothetical LOC254559	4,6	4,6E-58
L1CAM	L1 cell adhesion molecule	3,0	3,8E-56
ASPHD1	aspartate beta-hydroxylase domain containing 1	3,1	1,3E-55
SYN1	synapsin I	2,7	1,9E-49
RLTPR	RGD motif, leucine rich repeats, tropomodulin domain and proline-rich containing	1,9	1,6E-46
RUNDC3A	RUN domain containing 3A	3,8	2,5E-45
MMP24	matrix metalloproteinase 24 (membrane-inserted)	1,9	4,8E-42
SYP	synaptophysin	1,7	1,4E-39
KCNK3	potassium channel, subfamily K, member 3	4,6	1,0E-35
CELF3	trinucleotide repeat containing 4	3,3	1,6E-35
CACNA1B	calcium channel, voltage-dependent, N type, alpha 1B subunit	2,1	5,3E-34

C

REST-repressed genes in hypoxia (1 % oxygen)			
Gene symbol	Gene name	siREST-Fold change (log base 2)	p-Value
DISP2	dispatched homolog 2 (Drosophila)	1,96	3,5E-23
RAB3C	RAB3C, member RAS oncogene family	1,58	4,7E-08
IGSF21	immunoglobulin superfamily, member 21	1,01	5,5E-05
GANAB	glucosidase, alpha; neutral AB	0,93	3,7E-23
NES	nestin	0,91	2,3E-05
NR6A1	nuclear receptor subfamily 6, group A, member 1	0,85	8,5E-04
LBX1	ladybird homeobox 1	0,84	3,0E-03
DLX5	distal-less homeobox 5	0,82	5,3E-03
DUSP18	dual specificity phosphatase 18	0,72	1,5E-02
CYP1B1	cytochrome P450, family 1, subfamily B, polypeptide 1	0,70	1,2E-04
LOC100506433	hypothetical LOC100506433	0,69	4,9E-02
PAQR8	progesterin and adipoQ receptor family member VIII	0,66	9,5E-05
DOCK4	dedicator of cytokinesis 4	0,64	1,8E-03
HSDL2	hydroxysteroid dehydrogenase like 2	0,63	5,6E-08
MDGA2	MAM domain containing glycosylphosphatidylinositol anchor 2	0,62	1,7E-02
SCLY	selenocysteine lyase	0,61	1,4E-01
LAMA4	laminin, alpha 4	0,60	4,3E-04
GALNT14	UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 14 (GalNAc-T14)	0,60	1,6E-02
CHSY3	chondroitin sulfate synthase 3	0,57	1,9E-02
TMEM54	transmembrane protein 54	0,57	7,3E-04


D

REST repressed in normoxia and hypoxia




E





REST repressed in hypoxia only



F

Hypoxia Induced






B

Consensus RE1 motifs identified in the REST-dependent genes repressed in hypoxia

UCSC ID	Gene	RE1 position relative to the TSS	log2 odds score
uc003mtd.4	EXOC2	+637	17.36
uc021zal.1	PAQR8	-1968	15.26
uc004atz.3	FAM120A	-1886	15.4
uc001ksp.3	SFXN3	-421	20.74
uc001nua.4	GANAB	+145	15.49
uc001twy.4	TAOK3	+41	18.05
uc001wub.4	MIPOL1	-801	15.27
uc010ani.3	MDGA2	-817	17.36
uc002cks.3	CACNA1H	+681	16.8
uc010wtj.1	SEC14L1	-866	15.69
uc002wyf.3	DNMT3B	-1215	16.18
uc011ads.1	SYNJ1	-1549	20.59



Normoxic genes

0.7 %

(164 out of 23,284)

Hypoxic genes

20 %

(201 out of 1013)